

SLICC/ACR Damage Index (SDI)
**Systemic Lupus International Collaborating Clinics/American College of Rheumatology
Damage Index**

Damage occurring since diagnosis of SLE, ascertained by clinical assessment and present for ≥ 6 months unless otherwise stated. Repeat episodes have to be ≥ 6 months apart to score 2. The same lesion cannot be scored twice.

Note that you only need to check the box "1 point" when a descriptor is present; leave empty otherwise. Exceptions are descriptors that can generate scores of > 1 . These fields are required.

Ocular
Either eye, by clinical assessment

Any cataract ever 1 point

Retinal change or optic atrophy 1 point

Neuropsychiatric

Cognitive impairment (e.g. memory deficit, difficulty with calculation, poor concentration, difficulty in spoken or written language, impaired performance level) or major psychosis 1 point

Seizures requiring therapy for ≥ 6 months 1 point

Cerebral vascular accident ever (score 2 if > 1), or surgical resection for causes other than malignancy 0 points
 1 point
 2 points
 (Please note that this field is required)

Cranial or peripheral neuropathy (excluding optic) 1 point

Transverse myelitis 1 point

Renal

Estimated or measured GFR $< 50\%$ 1 point

Proteinuria ≥ 3.5 g/24 hours (score 1) or end-stage renal disease, regardless of dialysis or transplantation (score 3) 0 points
 1 point
 3 points
 (Please note that this field is required)

Pulmonary

Pulmonary hypertension (right ventricular prominence, or loud P2) 1 point

Pulmonary fibrosis (physical and X-ray) 1 point

Shrinking lung (X-ray) 1 point

Pleural fibrosis (X-ray) 1 point

Pulmonary infarction (X-ray) or surgical resection for causes other than malignancy 1 point

Cardiovascular

Angina or coronary artery bypass 1 point

Myocardial infarction ever (score 2 if > 1) 0 points
 1 point
 2 points
 (Please note that this field is required)

Cardiomyopathy (ventricular dysfunction) 1 point

Valvular disease (diastolic murmur, or a systolic murmur > 3/6) 1 point

Pericarditis (for ≥ 6 months) or pericardiectomy 1 point

Peripheral vascular

Claudication (for ≥ 6 months) 1 point

Minor tissue loss (pulp space) 1 point

Significant tissue loss ever (e.g. loss of digit or limb, resection) (score 2 if > 1) 0 points
 1 point
 2 points
 (Please note that this field is required)

Venous thrombosis with swelling, ulceration, or venous stasis 1 point

Gastrointestinal

Infarction or resection of bowel (below duodenum), spleen, liver or gall bladder ever (score 2 if > 1) 0 points
 1 point
 2 points
 (Please note that this field is required)

Mesenteric insufficiency 1 point

Chronic peritonitis 1 point

Stricture or upper gastrointestinal tract surgery ever 1 point

Pancreatic insufficiency requiring enzyme replacement or with pseudocyst 1 point

Musculoskeletal

Muscular atrophy or weakness 1 point

Deforming or erosive arthritis (including reducible deformities, excluding avascular necrosis) 1 point

Osteoporosis with fracture or vertebral collapse (excluding avascular necrosis) 1 point

Avascular necrosis (score 2 if > 1) 0 points
 1 point
 2 points
 (Please note that this field is required)

Osteomyelitis 1 point

Ruptured tendons 1 point

Skin

Scarring chronic alopecia 1 point

Skin ulceration (excluding thrombosis) for ≥ 6 months 1 point

Premature gonadal failure

Premature gonadal failure

1 point

Diabetes

Diabetes (regardless of treatment)

1 point

Malignancy

Malignancy (score 2 if > 1 site)
Exclude dysplasia.

- 0 points
 - 1 point
 - 2 points
- (Please note that this field is required)

Total SLICC/ACR Damage Index (SDI) score

Total SDI score
